

On the irreducibility of the two variable zeta-function for curves over finite fields

Niko Naumann

February 1, 2008

Abstract

In [P] R. Pellikaan introduced a two variable zeta-function $Z(t, u)$ for a curve over a finite field $\mathbb{F}_q$ which, for $u = q$, specializes to the usual zeta-function and he proved, among other things, rationality: $Z(t, u) = (1 - t)^{-1}(1 - ut)^{-1}P(t, u)$ with $P(t, u) \in \mathbb{Z}[t, u]$. We prove that $P(t, u)$ is absolutely irreducible. This is motivated by a question of J. Lagarias and E. Rains about an analogous two variable zeta-function for number fields.

MSC2000: 11G20 (primary), 14G10 (secondary)

1 Introduction

Let X be a proper, smooth, geometrically connected curve of genus g over the finite field $\mathbb{F}_q$. The zeta-function of $X/\mathbb{F}_q$ can be written as a power series

$$Z(t) = \sum_D \frac{q^{h^0(D)} - 1}{q - 1} t^{\deg(D)}.$$

Here the sum is over $\mathbb{F}_q$ -rational divisor classes of X and $h^0(D) := \dim_{\mathbb{F}_q} H^0(X, \mathcal{O}(D))$. Writing b_{nk} for the number of divisor classes

of degree n and with $h^0(D)=k$ this becomes

$$Z(t) = \sum_{n \geq 0} \sum_{k \geq 1} b_{nk} \frac{q^k - 1}{q - 1} t^n.$$

In [P] R. Pellikaan observed that the classical proof of rationality and the functional equation for $Z(t)$ go through when q is treated as a variable in this expression. He thus introduced the following power series in [P], Def. 3.1:

$$Z(t, u) := \sum_{n \geq 0} \sum_{k \geq 1} b_{nk} \frac{u^k - 1}{u - 1} t^n.$$

This is called the two variable zeta-function of the curve. We will denote by h the class-number of $X/\mathbb{F}_q$, i.e. $h = |\text{Pic}^0(X)|$. Then Pellikaan proved:

Theorem 1 *We have*

$$Z(t, u) = (1 - t)^{-1} (1 - ut)^{-1} P(t, u) \text{ with } P \in \mathbb{Z}[t, u].$$

Furthermore:

- 1) $\deg_t P = 2g$, $\deg_u P = g$.
- 2) In the expansion $P(t, u) = \sum_{i=0}^{2g} P_i(u) t^i$ one has $P_0(u) = 1$, $\deg_u P_i(u) \leq i/2 + 1$ and $P_{2g-i}(u) = u^{g-i} P_i(u)$ for $0 \leq i \leq 2g$.
- 3) $P(1, u) = h$.

Here $\deg_u$ and $\deg_t$ denote the degree of a polynomial in the indicated variable. The above results are all taken from [P], Prop. 3.5 and we copied only those needed later on. Note that the statement $\deg_u P_i(u) \leq i/2$ in [loc. cit.] is a misprint. Indeed, we will see below that one always has $\deg_u P_1(u) = 1$ (unless $g = 0$). As expected we have $P(t, u) = 1$ in case $g = 0$.

In [GS] G. van der Geer and R. Schoof used analogies from Arakelov-theory to define a two variable zeta-function for number fields along the above lines. As the number field case will serve only as a motivation in this note we refer to the original sources [GS] and [LR] for definitions and to [D] for a

comparison between them. Suffice it to say that in [LR], section 8 we find an entire function $\xi_{\mathbb{Q}}(w, s)$ of two complex variables which for $w = 1$ equals Riemann's ξ -function. In particular, the zeroes of $\xi_{\mathbb{Q}}(1, s)$ are precisely the non-trivial zeros of the Riemann zeta-function. One is thus led to study the zero-locus of $\xi_{\mathbb{Q}}(w, s)$. J. Lagarias and E. Rains ask whether it might be the closure of a single irreducible complex-analytic variety of multiplicity one. The corresponding question in the geometric case seems to be whether the zero-locus of $P(t, u)$ is irreducible.

This is indeed the case:

Theorem 2 *In the above situation, $P(t, u)$ is irreducible in $\mathbb{C}(u)[t]$.*

As an illustration we discuss the cases $g = 1$ and $g = 2$:

For $g = 1$ setting $N := |X(\mathbb{F}_q)|$ we have

$$P(t, u) = 1 + (N - 1 - u)t + ut^2,$$

c.f. [P], example 3.4. This polynomial is reducible in $\mathbb{C}(u)[t]$ if and only if $N = 0$ in which case we have $P(t, u) = (1 - t)(1 - ut)$. But it is well-known that a curve of genus one over a finite field always has a rational point, i.e. $N \neq 0$.

In case $g = 2$ let the usual zeta-function of X be

$$Z(t) = (1 - t)^{-1}(1 - qt)^{-1}L(t)$$

with

$$L(t) = 1 + at + bt^2 + qat^3 + q^2t^4$$

for certain $a, b \in \mathbb{Z}$. As X is hyperelliptic, Prop. 4.3. of [P] can be used to compute

$$P(t, u) = 1 + ((a+q)-u)t + ((q(q-1)+aq+b)-(a+q-1)u)t^2 + ((a+q)-u)ut^3 + u^2t^4.$$

This will not be used in the sequel and we omit the proof.

In order not to lead intuition astray we point out that in general $Z(t, u)$ is not determined by $Z(t)$, see [P], example 4.4.

After a lengthy computation with discriminants one sees that a necessary condition for this $P(t, u)$ to be reducible is

$$b + a(q + 1) + (q^2 + 1) = 0.$$

But this expression equals $L(1) = h \neq 0$!

The fact that $h \neq 0$ enters in the general proof of theorem 2 precisely through the condition $\beta \neq 0$ of lemma 3 below. Note, however, that condition 2) of this lemma cannot be dropped. So one needs one more result on $P(t, u)$, contained in Prop. 4, which follows from Clifford's theorem.

I would like to thank C. Deninger for posing the above problem and for useful discussions on the topic.

2 Proof

We will use the following criterion for irreducibility:

Lemma 3 *Let k be a field, $F \in k[u, t]$ and assume:*

- 1) *F is monic in t .*
- 2) *the leading coefficient of F as a polynomial in u is irreducible in $k[t]$.*
- 3) *there are $\alpha, \beta \in k$, $\beta \neq 0$ with $F(u, \alpha) = \beta$.*

Then F is irreducible in $k(u)[t]$.

This lemma will be applied to

$$F = \tilde{P}(t, u) := t^{2g}P(t^{-1}, u) \in \mathbb{C}[u, t].$$

Note that the irreducibility of $\tilde{P}$ in $\mathbb{C}(u)[t]$ will imply the irreducibility of P because $P(0, u) = P_0(u) = 1 \neq 0$ by theorem 1. The advantage of $\tilde{P}$ is that it is monic in t and so satisfies condition 1) of lemma 3. Also 3) is satisfied (with $\alpha = 1$, $\beta = h$) according to theorem 1, 3).

Proof (of lemma 3) Assume to the contrary that $F = fg$ in $k(u)[t]$ with f and g of positive degree and monic. One knows, c.f. for example [E], Prop 4.11, that the coefficient of f and g are integral over $k[u]$ and as $k[u]$ is integrally closed we have $f, g \in k[u, t]$. So we can consider the decomposition $F = fg$ as polynomials in u and infer from 2) that the leading coefficient of f as a polynomial in u lies in $k[t]^* = k^*$ (upon exchanging f and g if necessary). In particular $n := \deg_u f(u, t) = \deg_u f(u, \alpha)$. Substituting 3)

gives $\beta = f(u, \alpha)g(u, \alpha)$ in $k[u]$. As $\beta \neq 0$ we get $n = 0$, i.e. f is constant in u hence $f \in k^*$, contradiction.

□

We are left with verifying condition 2) of lemma 3 for the given $\tilde{P}$, i.e. the leading coefficient of $\tilde{P}$ as a polynomial in u is irreducible in $k[t]$. We will in fact determine this coefficient:

Proposition 4 *For $g \geq 1$: $\tilde{P}(t, u) = (1 - t)u^g + O(u^{g-1})$.*

Proof We already know $\deg_u \tilde{P} = g$. Also the assertion is clear for $g = 1$ from the formula for $P(t, u)$ recalled in the introduction. We assume $g \geq 2$. Looking at

$$\tilde{P}(t, u) = t^{2g} + P_1(u)t^{2g-1} + \cdots + P_g(u)t^g + uP_{g-1}(u)t^{g-1} + \cdots + u^gt^0$$

and using the bound $\deg_u P_i(u) \leq i/2 + 1$ we see that u^g can only occur in the last three terms: $u^{g-2}P_2(u)t^2 + u^{g-1}P_1(u)t + u^g$. So the proof is completed by the following result on P_1 and P_2 .

□

Proposition 5 *For $g \geq 1$:*

- 1) $\deg_u P_1(u) = 1$ and the leading coefficient is -1 .
- 2) $\deg_u P_2(u) \leq 1$.

Proof This is again clear for $g = 1$. We assume $g \geq 2$ and write $P_i(u) = \sum_k \alpha_{ik}u^k$, $\alpha_{ik} \in \mathbb{Z}$. As we already know $\deg_u P_1(u) \leq 1$ and $\deg_u P_2(u) \leq 2$ we need to show $\alpha_{11} = -1$ and $\alpha_{22} = 0$. Recalling the notation b_{nk} from the introduction we have the following

Claim: $b_{12} = \alpha_{00} + \alpha_{11}$ and $b_{23} = \alpha_{00} + \alpha_{11} + \alpha_{22}$.

Granting this we observe that Clifford's theorem, c.f. [H], IV, thm 5.4, gives $b_{12} = b_{23} = 0$. Recalling also $\alpha_{00} = 1$, because $P_0(u) = 1$, and substituting gives indeed $\alpha_{11} = -1$ and $\alpha_{22} = 0$.

□

To prove the above claim we write the rational expression for $Z(t, u)$ in theorem 1 in terms of coefficients:

$$Z(t, u) = \sum_{n \geq 0} \sum_{k \geq 1} b_{nk} \frac{u^k - 1}{u - 1} t^n = \left(\sum_{i \geq 0} t^i \right) \left(\sum_{j \geq 0} (ut)^j \right) \left(\sum_{l \geq 0} \sum_{k \geq 0} \alpha_{lk} u^k t^l \right).$$

This gives

Lemma 6

- 1) for $\nu, \alpha \geq 0$: $\sum_{k \geq \alpha+1} b_{\nu k} = \sum_{\mu, i \geq 0, \mu+i \leq \nu} \alpha_{i, \alpha-\mu}$
- 2) for $\nu \geq 0, \mu \geq 1$: $b_{\nu \mu} = \sum_{i=0}^{\nu} (\alpha_{i, \mu-\nu-1+i} - \alpha_{i \mu})$

We omit the details of this straightforward computation except to say that for 1) one uses $\frac{u^k-1}{u-1} = 1 + \dots + u^{k-1}$ and 2) follows from 1) by a telescope-summation. In the formulation of the lemma it is understood that $\alpha_{nk} = 0$ whenever $k < 0$. We get from 2):

$$b_{23} = \alpha_{00} - \alpha_{03} + \alpha_{11} - \alpha_{13} + \alpha_{22} - \alpha_{23}$$

and

$$b_{12} = \alpha_{00} - \alpha_{02} + \alpha_{11} - \alpha_{12}.$$

But we know $\alpha_{02} = \alpha_{03} = \alpha_{12} = \alpha_{13} = \alpha_{23} = 0$ because $\deg_u P_i(u) \leq i/2 + 1$. This concludes the proof of the claim, hence of theorem 2.

References

- [D] C. Deninger, A regularized product expansion for the two-variable zeta-function of van der Geer and Schoof, preprint, 2002.
- [E] D. Eisenbud, Commutative Algebra with a View Towards Algebraic Geometry, GTM 150, Springer, 1995.
- [GS] G. van der Geer, R. Schoof, Effectivity of Arakelov Divisors and the Theta Divisor of a Number Field, Selecta Math. (N.S.), **6**, 2000, no. 4, pp 377-398.

- [H] R. Hartshorne, Algebraic Geometry, GTM 52, Springer, 1977.
- [LR] J. Lagarias, E. Rains, On a Two-Variable Zeta Function for Number Fields, arXiv:math.NT/0104176 v5 7 Jul 2002.
- [P] R. Pellikaan, On special divisors and the two variable zeta function of algebraic curves over finite fields, in: Arithmetic, Geometry and Coding Theory 4, Luminy, 1993, Walter de Gruyter, Berlin, 1996, pp 175-184.

Mathematisches Institut der WWU Münster
Einsteinstr. 62
48149 Münster
Germany
e-mail: naumannn@uni-muenster.de